

COT 3420  
SUMMER A 2003  
Section 1

EXAM # 1 ANSWERS

**Question 1.**(20 points)

1. d    2. c    3. b    4. b    5. b    6. c    7. c    8. a    9. a    10. c

**Question 2.** (20 points)

The proof is by structural induction on  $F$ .

Case 1:  $F$  is an atom. Then  $F = P_i$  for some  $i \in N$ . So,  $n[atom, P_i] = 1$ , and  $n[con, P_i] = 0$ . The equation  $n[atom, F] = n[con, P_i] + 1$  becomes

$$1 = 0 + 1$$

which is true.

Case 2:  $F = \neg G$ . We assume that the equation holds for  $G$ .

$$(IH) \quad n[atom, G] = n[con, G] + 1$$

Now we relate the  $F$  counts to the  $G$  counts.

$$(1) \quad n[atom, F] = n[atom, G] \quad \text{because } F = \neg G \text{ and } \neg \text{ is not an atom}$$

$$(2) \quad n[con, F] = n[con, G] \quad \text{because } F = \neg G \text{ and } \neg \text{ is not a binary connective}$$

Now we compute  $n[atom, F]$ .

$$n[atom, F] = n[atom, G] \quad \text{by (1)}$$

$$= n[con, G] + 1 \quad \text{by (IH)}$$

$$= n[con, F] + 1 \quad \text{by (2)}$$

So,  $n[atom, F] = n[con, F] + 1$ .

Cases 3, 4, 5, 6:  $F = (GCH)$ . We assume that the proposition is true for  $G$  and  $H$ .

$$(IH1) \quad n[atom, G] = n[con, G] + 1$$

$$(IH2) \quad n[atom, H] = n[con, H] + 1$$

We express the  $F$  counts as functions of  $G$  and  $H$  counts. Since  $F = (GCH)$  and the symbols  $(, C, )$  are not atoms,

$$(3) \quad n[atom, F] = n[atom, G] + n[atom, H].$$

The connectives of  $F$  are the connectives of  $G$  and  $H$  plus  $C$ .

$$(4) \quad n[con, F] = n[con, G] + 1 + n[con, H]$$

Now we relate  $n[atom, F]$  to  $n[con, F]$ .

$$\begin{aligned}
n[atom, F] &= n[atom, G] + n[atom, H] && \text{by (3)} \\
&= n[con, G] + 1 + n[atom, H] && \text{by (IH1)} \\
&= n[con, G] + 1 + n[con, H] + 1 && \text{by (IH2)} \\
&= (n[con, G] + 1 + n[con, H]) + 1 && \text{by grouping} \\
&= n[con, F] + 1 && \text{by (4)}
\end{aligned}$$

So,  $n[atom, F] = n[con, F] + 1$ .

Grading Criteria: 1. Listing the 6 cases: 2 points.

2. Case 1: 2 points.

3. Case 2: 7 points; The IH - 1 point, the 2 count equations - 2 points, the derivation - 3 points, the reasons - 1 point.

4. Cases 3,4,5,6: 9 points; IH's - 2points, the count equations - 2 points, the derivation - 4 points, the reasons - 1 point.

**Question 3.** (10 points)

The tree is shown in Figure 1.

Grading Criteria: 1 point off for each out of order node.

**Question 4.** (15 points)

The proof is by mathematical induction on  $n$ .

Basis:  $n = 0$ . Then  $\bigvee_{i=0}^0 P_i = P_0$ , a symbol. So,  $|\bigvee_{i=0}^0 P_i| = 1$ , and the equation  $|\bigvee_{i=0}^n P_i| = 4n + 1$  becomes

$$1 = 4 * 0 + 1$$

which is true.

Inductive Step: Assume that the equation is true for  $n$ .

$$(IH) |\bigvee_{i=0}^n P_i| = 4n + 1$$

Let us prove

$$|\bigvee_{i=0}^{n+1} P_i| = 4(n + 1) + 1$$

The know that  $\bigvee_{i=0}^{n+1} P_i$  denotes the string  $(\bigvee_{i=0}^n P_i \vee P_{n+1})$ .

So, the length of  $\bigvee_{i=0}^{n+1} P_i$  is the length of  $\bigvee_{i=0}^n P_i$  plus the lengths of the symbols  $(, \vee, P_{n+1}$  and  $)$ . So,

$$(1) |\bigvee_{i=0}^{n+1} P_i| = |\bigvee_{i=0}^n P_i| + 4$$

Now we relate  $|\bigvee_{i=0}^{n+1} P_i|$  to  $n$ .

$$|\bigvee_{i=0}^{n+1} P_i| = |\bigvee_{i=0}^n P_i| + 4 \quad \text{by (1)}$$

$$= (4n + 1) + 4 \quad \text{by (IH)}$$

$$= (4n + 4) + 1 \quad \text{by grouping}$$

$$= 4(n + 1) + 1 \quad \text{by distributivity}$$

So, we proved that  $|\bigvee_{i=0}^{n+1} P_i| = 4(n + 1) + 1$ .

Grading Criteria: 1. Listing the 2 cases: 2 points.




Figure 1: The answer for Question 3

2. Basis: 3 points.
3. Inductive Step: 10 points. The IH - 1 point; the formula (1) - 5 points, the derivation + reasons=4 points.

**Question 5.** (10 points)

$$S = \{P_1, P_2, P_3, P_0, P_4, P_7, \neg P_2, \neg P_3, (P_4 \longleftrightarrow P_7), (P_1 \vee \neg P_2), (P_2 \wedge \neg P_3), (P_0 \wedge (P_4 \longleftrightarrow P_7)), ((P_1 \vee \neg P_2) \longrightarrow (P_2 \wedge \neg P_3)), \neg(P_0 \wedge (P_4 \longleftrightarrow P_7)), \neg((P_1 \vee \neg P_2) \longrightarrow (P_2 \wedge \neg P_3)), (\neg((P_1 \vee \neg P_2) \longrightarrow (P_2 \wedge \neg P_3)) \longleftrightarrow \neg(P_0 \wedge (P_4 \longleftrightarrow P_7)))\}$$

Grading Criteria:  $\frac{10}{16}(\text{number-of-right-answers} - \text{number-of-wrong-answers})$